

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SUPT. HARD'S NEW BOOK ON MUSHROOMS

AN ADVANCE REVIEW BY W. A. KELLERMANN.

Now comes forth an announcement of a new book on Mushrooms by that mushroom-hunter, mushroom-eater, mushroom-writer, as it were, mushroom-fiend—but we have seen the MSS., and the pictures, and can quiet the mushroom public by saying that the book is all right. In natural history no one is "authority"—each one thinks for himself, speaks for himself; it is the plan we here adopt to give a single description of Mr. Hard's, and a few sample illustrations, which his publishers have kindly allowed me to use in advance. We take at random a description, say of AMANITA MAPPA, and it will appear as follows in the new book:

"AMANITA MAPPA FR. THE DELICATE AMANITA. Poisonous.

"Mappa means a napkin; so-called from the volva.

"The pileus is two to three inches broad, convex, then expanded, plane, obtuse or depressed, without separable cuticle; margin nearly even; white or yellowish, usually with patches of the volva, dry.

"The gills are adnexed, close, narrow, shining, white.

"The stem is two to three inches long, stuffed, then hollow, cylindrical, nearly smooth, bulbous, nearly globose at the base, white, almost equal above the bulb.

"The volva with its free margin is acute and narrow.

ring is membranaceous, superior, soft, lax, ragged.
"Its color is quite as variable and its habits are much like A. phalloides, from which it can only be distinguished by its less developed volva which, instead of being cup-shaped, is little more than a mere rim fringing the bulb. The odor at times is very strong. It is found in open woods and under brush. Label it poisonous.'

Those interested to know more about this publication, Mushrooms Edible and Otherwise, and to place orders at once, are referred to the author (Supt. M. E. Hard, Chillicothe, O.), but it may be added that it will be very comprehensive, describing and figuring practically all the mushrooms any one can find in this country; also explaining the structure of the plants, the descriptive terms used, classification; giving keys and directions for study, a glossary, recipes for cooking, and a chapter on Culture of Mushrooms.

The figures we use in this number of the BULLETIN, taken from the book now in press, are as follows:

Fig. 277. AMANITA MAPPA. Poisonous. Natural size. See description quoted above.

Fig. 278. COLLYBIA CONFLUENS. Edible. Natural size. Growing among leaves in the woods after warm rains, in tufts, sometimes in rows or lines. The cap is an inch to an inch and a quarter broad, reddish brown; stems two to three inches long, hollow, pale red, sprinkled with a mealy pubescence.

Fig. 279. CLAVARIA SPINULOSA. Edible. Color somewhat cinnamon brown throughout. In mixed woods, after frequent

rains in August to October.

Fig. 280. TRICHOLOMA MACULATESCENS. The spotted Tricholoma, so-called because the cap in drying becomes more or less spotted. The cap is one ane a half to three inches broad, reddish-brown. Does not seem to be very common.

Fig. 281. Cantharellus floccosus. The Woolly Cantharellus. Edible. The cut from a photograph by C. G. Lloyd. It is elongated, funnel-shaped or trumpet-shaped, ochraceous yellow, one to two inches broad at the top, and never more than four inches high.



Fig. 277. $A_{\rm M-AN-I'-TA}$ MAP'-PA. From Hard's new book on Mushrooms.




Fig. 278. Col.-Lyb'-1-A con'-FLU-ENS. From Hard's new book on Mushrooms.





Fig. 279. CLA-VA'-RI-A SPIN-U-LO'-SA. From Hard's new book on Mushrooms.



Fig. 280. Tri-cho-lo'-ma mac-u-la-tes'-cens. From Hard's new book on Mushrooms.



The Mycological Bulletin is issued Monthly, Price 25c. Copies of Vol. II (1904), Vol. III (1905), Vol. IV (1906) and Vol. V (1907) may be had for 50 cents each, or cloth bound copies for 75 cents. No copies remain of Vol. I (1902). Address W. A. Kellerman, Columbus, Ohio.

Fig. 281. CAN-THA-REL'-LUS FLOC-CO'-SUS. From Hard's new bookon Mushrooms.